

TEMPORAL 1-SOLITON SOLUTION OF THE COMPLEX GINZBURG-LANDAU EQUATION WITH POWER LAW NONLINEARITY

A. Biswas

Center for Research and Education in Optical Sciences
and Applications
Department of Mathematical Sciences
Delaware State University
Dover, DE 19901-2277, USA

Abstract—This paper obtains the exact 1-soliton solution of the complex Ginzburg-Landau equation with power law nonlinearity that governs the propagation of solitons through nonlinear optical fibers. The technique that is used to carry out the integration of this equation is He's semi-inverse variational principle.

1. INTRODUCTION

The study of the propagation of solitons through optical fibers has been going on for the past few decades [1–25]. The nonlinear Schrödinger's equation (NLSE) is the main equation that studies this process [1]. There are various variants of this equation that also models this physical phenomena depending on the nonlinear perturbation effects. One such equation is the Radhakrishnan, Kundu, Lakshmanan (RKL) equation [22]. In this paper, the complex Ginzburg-Landau (CGL) equation, with power law nonlinearity, will be studied, which models optical solitons with a few perturbation effects.

In the presence of perturbation effects, the adiabatic parameter dynamics of optical solitons can be obtained also by using various mathematical techniques. They are Variational Principle [1], Soliton Perturbation Theory [9], Collective Variables Method [16] and others. But as a matter of fact, none of these methods can integrate the CGL equation. There has been a newly developed method called the He's Variational Principle (HVP) [10] that can carry out the integration of CGL equation. In this paper, HVP will be employed to carry out

Corresponding author: A. Biswas (biswas.anjan@gmail.com).

the integration of the CGL equation with power law nonlinearity. The closed form 1-soliton solution will also enable to obtain the parameter restrictions for the solitons to exist.

2. MATHEMATICAL ANALYSIS

The dimensionless form of the CGL equation, with power law nonlinearity, which will be studied in this paper, is given by [17]

$$iq_t + aq_{xx} + b(|q|^{2m})q = \alpha \frac{|q_x|^2}{q^*} + \frac{\beta}{4|q|^2 q^*} \left[2|q|^2 (|q|^2)_{xx} - \left\{ (|q|^2)_x \right\}^2 \right] + i\gamma q \quad (1)$$

where x represents the non-dimensional distance along the fiber while; t represents time in dimensionless form; a , b , α , β and γ are real valued constants. The coefficients of a and b are due to dispersion and power law nonlinearity where the parameter m dictates the power law nonlinearity. The terms due to α , β and γ are from the perturbation effects [10, 17].

Equation (1) is a nonlinear partial differential equation that is not integrable by the classical method of Inverse Scattering Transform since (1) will fail the Painleve test of integrability [1]. However, the HVP will be available for integrating (1), and therefore this technique will be used in this paper to carry out the integration of (1).

The starting point is the hypothesis [10]:

$$q(x, t) = g(x - vt)e^{i(-\kappa x + \omega t + \theta)} \quad (2)$$

where the function g represents the pulse shape, and v is the velocity of the soliton. From the phase component, κ is the soliton frequency; ω is the soliton wave number, while θ is the phase constant. Substituting this hypothesis into (1) and decomposing into real and imaginary parts respectively yield

$$-\omega g + a(g'' - \kappa^2 g) + bg^{2m+1} = \frac{\alpha}{g} \left\{ (g')^2 + \kappa^2 g^2 \right\} + \beta g'' \quad (3)$$

and

$$g'(v + 2a\kappa) = -\gamma g \quad (4)$$

where $g' = dg/ds$ and $g'' = d^2g/ds^2$ with

$$s = x - vt \quad (5)$$

Now (4) gives the velocity of the soliton as

$$v = -2a\kappa - \frac{\ln g}{t} \quad (6)$$

Multiplying (3) by g' and integrating, yields

$$-\left\{\omega + \kappa^2(a + \alpha) + \frac{\alpha\gamma^2}{(v + 2a\kappa)^2}\right\} \frac{g^2}{2} + (\alpha - \beta) \frac{g'^2}{2} + \frac{bg^{2m+2}}{2m+2} = K \quad (7)$$

where K is the constant of integration. At this point, the quantity J is defined as

$$J = \int_{-\infty}^{\infty} K ds = \int_{-\infty}^{\infty} \left[-\left\{\omega + \kappa^2(a + \alpha) + \frac{\alpha\gamma^2}{(v + 2a\kappa)^2}\right\} \frac{g^2}{2} + (\alpha - \beta) \frac{g'^2}{2} + \frac{bg^{2m+2}}{2m+2} \right] ds \quad (8)$$

Now, the 1-soliton solution ansatz, given by [10]

$$g(s) = \frac{A}{\cosh^{\frac{1}{m}}(Bs)} \quad (9)$$

is substituted into (8). Here, in (9), the parameters A and B represent the amplitude and inverse width of the soliton respectively. He's semi-inverse variational principle states that the parameters A and B are determined from the solution of the equations [10]

$$\frac{\partial J}{\partial A} = 0 \quad (10)$$

and

$$\frac{\partial J}{\partial B} = 0. \quad (11)$$

Thus, by virtue of the soliton hypothesis given by (9), Equation (8) reduces to

$$J = \left[-\frac{A^2}{2B} \left\{ \omega + \kappa^2(a + \alpha) + \frac{\alpha\gamma^2}{(v + 2a\kappa)^2} \right\} + \frac{m(\alpha - \beta)A^2B}{2(m+2)} + \frac{bA^{2m+2}}{(m+1)(m+2)B} \right] \frac{\Gamma(\frac{1}{2})\Gamma(\frac{1}{m})}{\Gamma(\frac{1}{m} + \frac{1}{2})} \quad (12)$$

From (12), Equations (10) and (11) respectively reduce to

$$\frac{m(\alpha - \beta)B^2}{m+2} + \frac{2bA^{2m}}{m+2} = \omega + \kappa^2(a + \alpha) + \frac{\alpha\gamma^2}{(v + 2a\kappa)^2} \quad (13)$$

and

$$\frac{m(\alpha - \beta)B^2}{m+2} - \frac{2bA^{2m}}{(m+1)(m+2)} = -\left\{ \omega + \kappa^2(a + \alpha) + \frac{\alpha\gamma^2}{(v + 2a\kappa)^2} \right\} \quad (14)$$

After solving (13) and (14) yields

$$A = \left[\frac{m+1}{b(v+2a\kappa)^2} \{ \alpha\gamma^2 + \omega(v+2a\kappa)^2 + \kappa^2(a+\alpha)(v+2a\kappa)^2 \} \right]^{\frac{1}{2m}} \quad (15)$$

and

$$B = \frac{1}{v+2a\kappa} \left[\frac{\alpha\gamma^2 + \omega(v+2a\kappa)^2 + \kappa^2(a+\alpha)(v+2a\kappa)^2}{\beta - \alpha} \right]^{\frac{1}{2}} \quad (16)$$

Also from (13) and (14), one can obtain the relation between the soliton amplitude A and inverse width B as

$$B = A^m \sqrt{\frac{b}{(m+1)(\beta - \alpha)}} \quad (17)$$

Now, since A and B are constants, Equation (17) also implies that

$$\frac{b}{\beta - \alpha} = \text{constant} > 0 \quad (18)$$

Also, (17) yields the domain restriction as

$$b(\beta - \alpha) > 0 \quad (19)$$

Finally, Equations (15) and (16) respectively imply

$$b \{ \alpha\gamma^2 + \omega(v+2a\kappa)^2 + \kappa^2(a+\alpha)(v+2a\kappa)^2 \} > 0 \quad (20)$$

and

$$(\beta - \alpha) \{ \alpha\gamma^2 + \omega(v+2a\kappa)^2 + \kappa^2(a+\alpha)(v+2a\kappa)^2 \} > 0 \quad (21)$$

Thus, finally the 1-soliton solution to (1) is given by

$$q(x, t) = \frac{A}{\cosh^{\frac{1}{m}} [B(x - vt)]} e^{i(-\kappa x + \omega t + \theta)} \quad (22)$$

where the amplitude and width are given by (15) and (16), and the velocity of the soliton is given by (6) while the constraint relation between the soliton parameters are given by (18)–(21).

3. CONCLUSION

This paper obtains the 1-soliton solution of the CGL equation with power law nonlinearity. the HVP is used to carry out the integration. In the process of determining the solution, several constraint relations between the soliton parameters are obtained. In future, this technique can be applied to CGL equation with perturbation terms.

ACKNOWLEDGMENT

This research work was fully supported by NSF-CREST Grant No: HRD-0630388 and Army Research Office (ARO) along with the Air Force Office of Scientific Research (AFOSR) under the award number: W54428-RT-ISP and these supports are genuinely and sincerely appreciated.

REFERENCES

1. Biswas, A. and S. Konar, *Introduction to Non-kerr Law Optical Solitons*, CRC Press, Boca Raton, FL, USA, 2006.
2. Biswas, A., "Dynamics of Gaussian and super-Gaussian solitons in birefringent optical fibers," *Progress In Electromagnetics Research*, Vol. 33, 119–139, 2001.
3. Biswas, A. and S. Konar, "Theory of dispersion-managed optical solitons," *Progress In Electromagnetics Research*, Vol. 50, 83–134, 2005.
4. Biswas, A., "Stochastic perturbation of parabolic law optical solitons," *Journal of Electromagnetic Waves and Applications*, Vol. 21, No. 11, 1479–1488, 2007.
5. Biswas, A., D. Milovic, F. Majid, and R. Kohl, "Optical soliton sootling in a saturable law media," *Journal of Electromagnetic Waves and Applications*, Vol. 22, 1735–1746, 2008.
6. Biswas, A., R. Kohl, M. E. Edwards, and E. Zerrad, "Soliton parameter dynamics in a non-Kerr law media," *Progress In Electromagnetics Research C*, Vol. 1, 1–35, 2008.
7. Khalique, C. M. and A. Biswas, "Optical solitons with parabolic and dual-power law nonlinearity via Lie symmetry analysis," *Journal of Electromagnetic Waves and Applications*, Vol. 23, No. 7, 963–973, 2009.
8. Kohl, R., A. Biswas, D. Milovic, and E. Zerrad, "Adiabatic dynamics of Gaussian and super-Gaussian solitons in Dispersion-Managed optical fibers," *Progress In Electromagnetics Research*, PIER 84, 27–53, 2008.
9. Kohl, R., A. Biswas, D. Milovic, and E. Zerrad, "Soliton perturbation theory for dispersion-managed optical fibers," *Journal of Nonlinear Optical Physics and Materials*, Vol. 18, No. 2, 227–270, 2009.
10. Kohl, R., A. Biswas, D. Milovic, and E. Zerrad, "Optical solitons by He's variational principle in a non-Kerr law media," *Journal of*

- Infrared, Millimeter and Terahertz Waves*, Vol. 30, No. 5, 526–537, 2009.
11. Lott, D. A., A. Henriquez, B. J. M. Sturdevant, and A. Biswas, “A numerical study of optical soliton-like structures resulting from the nonlinear Schrödinger’s equation with square-root law nonlinearity,” *Applied Mathematics and Computation*, Vol. 207, No. 2, 319–326, 2009.
 12. Mishra, M. and S. Konar, “Interaction of solitons in a dispersion-managed optical communication system with asymmetric dispersion map,” *Journal of Electromagnetic Waves and Applications*, Vol. 21, No. 4, 2049–2058, 2007.
 13. Mishra, M. and S. Konar, “High bit rate dense dispersion-managed optical communication systems with distributed amplification,” *Progress In Electromagnetics Research*, PIER 78, 301–320, 2008.
 14. Panajotovic, A., D. Milovic, and A. Biswas, “Influence of even order dispersion on soliton transmission quality with coherent interference,” *Progress In Electromagnetics Research B*, Vol. 3, 63–72, 2008.
 15. Shwetanshumala, S., S. Jana, and S. Konar, “Propagation of a mixture of modes of a laser beam in a medium with saturable nonlinearity,” *Journal of Electromagnetic Waves and Applications*, Vol. 20, No. 1, 65–77, 2006.
 16. Shwetanshumala, S. and A. Biswas, “Femtosecond pulse propagation in optical fibers under higher order effects: A collective variables approach,” *International Journal of Theoretical Physics*, Vol. 47, No. 6, 1699–1708, 2008.
 17. Shwetanshumala, S., “Temporal solitons of modified complex Ginzburg-Landau equation,” *Progress In Electromagnetic Research Letters*, Vol. 3, 17–24, 2008.
 18. Srivastava, S., “Coupled bright-dark soliton pairs in polymeric photorefractive materials,” *Journal of Electromagnetic Waves and Applications*, Vol. 22, No. 2–3, 335–342, 2008.
 19. Stojanovic, B., D. Milovic, and A. Biswas, “Timing shift of optical pulses due to inter-channel cross-talk,” *Progress In Electromagnetic Research M*, Vol. 1, 21–30, 2008.
 20. Sturdevant, B. J. M., D. A. Lott, and A. Biswas, “Dynamics of topological optical solitons with time-dependent dispersion, nonlinearity and attenuation,” *Communications in Nonlinear Science and Numerical Simulation*, Vol. 14, No. 8, 3305–3308, 2009.

21. Sturdevant, B., D. A. Lott, and A. Biswas, "Topological solitons in 1+2 dimensions with time-dependent coefficients," *Progress In Electromagnetic Research Letters*, Vol. 10, 69–75, 2009.
22. Sturdevant, B., D. A. Lott, and A. Biswas, "Dark 1-soliton solution of the generalized Radhakrishnan, Kundu, Lakshmanan equation with nonlinear dispersion," Submitted.
23. Wazwaz, A. M., "Reliable analysis for nonlinear Schrödinger equations with a cubic nonlinearity and a power law nonlinearity," *Mathematical and Computer Modelling*, Vol. 43, No. 1–2, 178–184, 2006.
24. Wazwaz, A. M., "Exact solutions for the fourth order nonlinear Schrödinger equations with cubic and power law nonlinearities," *Mathematical and Computer Modelling*, Vol. 43, No. 7–8, 802–808, 2006.
25. Wazwaz, A. M., "A study on linear and nonlinear Schrödinger equations by the variational iteration method," *Chaos, Solitons and Fractals*, Vol. 37, No. 4, 1136–1142, 2008.